

Exact results for one dimensional stochastic cellular automata for different types of updates

N. Rajewsky¹ and M. Schreckenberg²

¹ Institut für Theoretische Physik
Universität zu Köln
D-50937 Köln, F.R.G.

² FB 11-Mathematik
Gerhard-Mercator-Universität Duisburg
D-47048 Duisburg, F.R.G.

Abstract We study two common types of time-noncontinuous updates for one dimensional stochastic cellular automata with arbitrary nearest neighbor interactions and arbitrary open boundary conditions. We first construct the stationary states using the matrix product formalism. This construction then allows to prove a general connection between the stationary states which are produced by the two different types of updates.

Using this connection, we derive explicit relations between the densities and correlation functions for these different stationary states.

¹email: nr@thp.uni-koeln.de

²email: schreck@math.uni-duisburg.de

In this letter we study one dimensional stochastic cellular automata with different types of update. We consider a chain of sites $i = 1, \dots, L$ (L even) and assign a discrete variable $\tau_i = 1, \dots, m$ to each site i . τ_i can for example represent m different sorts of particles. We then prescribe some local rule $\mathcal{T}$ describing the interaction between a pair of nearest neighbor variables (τ_i, τ_{i+1}) and, as usually done, choose this rule to be independent of the location of the sites. One can further allow for some rules $\mathcal{R}$ and $\mathcal{L}$, acting on the sites at the two ends of the chains. Such boundary conditions could for example mimic the injection and removal of particles with certain rates. This type of boundaries can lead to a rich dynamical behaviour of the model. One prominent example is the asymmetric exclusion model (see [1], [2] and ref. therein) which exhibits boundary induced phase transitions where the bulk density of particles depends in a subtle way on the boundary conditions. There are now several ways to implement our prescribed update rules. These implementations are an essential part of the definition of a model since they generally produce different transients, stationary states and averages of observables.

For analytic calculations one usually uses the *random sequential* update. One picks at random site i and applies the update rule. This update generally produces the weakest correlations between the sites compared to other update procedures. The master equation which describes the temporal evolution of the weights of the configurations $\{\tau_1, \tau_2, \dots, \tau_L\}$ can then be written as a Schrödinger-like equation with a (non-hermetian) hamiltonian consisting of a sum of local hamiltonians. For this update it was recently shown, that the stationary state can be always written as the scalar product of the product of some (generally) noncommuting matrices [9].

There are three other basic ways of implementing the update rules which are much more common for practical purposes like computer simulations, since the random sequential update requires an additional random number for each local update. These three types are called *sequential*, *sublattice parallel* and *parallel* update. In the parallel update one applies the update rules simultaneously to all sites. This case will not be discussed in the following. Note that the parallel update usually produces the strongest correlations between the sites and is often used in the context of traffic flow models [4].

In the sequential update we start at the right end of the chain and apply our boundary condition $\mathcal{R}$ to site $i = L$. We then update the pair (τ_{L-1}, τ_L) . After that we proceed with the pair (τ_{L-2}, τ_{L-1}) and so forth until we reach the left end of the chain where we apply $\mathcal{L}$ at site $i = 1$. This procedure can of course also be done from the left to the right. Let us denote these two updates with $T_{\rightarrow}$ and $T_{\leftarrow}$.

The sublattice parallel update $T_{\parallel}$ is defined as follows: In the first time step

one applies $\mathcal{L}$, $\mathcal{R}$ and updates all pairs (τ_i, τ_{i+1}) with an even i . In the second time step one updates all pairs (τ_i, τ_{i+1}) with an odd i .

In the following, we show that in the stationary state averages like densities and correlation functions of the sequential and sublattice parallel update are essentially the same. This is a nontrivial result since the microscopic configurations of the system are generally very different after applying a sequential or a sublattice parallel update. This result should be of practical use for people doing computer simulations of cellular automata or reaction-diffusion processes. Note that the sequential update can transport a particle many sites during one update of the whole chain while $T_{\parallel}$ can move a particle only two sites at maximum.

We start with the construction of the stationary state of an arbitrary model with *sequential* update $T_{\leftarrow}$ from the right to the left end. We will first show that it is always possible to write the stationary probability distribution $P_0(\tau_1, \tau_2, \dots, \tau_L)$ of the configurations $\{\tau_i\}$ as

$$P_0(\tau_1, \tau_2, \dots, \tau_L)_{\leftarrow} = \langle W | A_{\tau_1} A_{\tau_2} \dots A_{\tau_L} | V \rangle \quad (1)$$

where the A_{τ_i} are matrices (representating single site states), the $\langle W |, |V \rangle$ vectors (reflecting the influence of the boundary conditions) acting in some auxiliary space. One can formally define a column vector $A = (A_1, \dots, A_m)$ and rewrite (1) as

$$|P_0\rangle_{\leftarrow} = \langle W | A^{\otimes L} | V \rangle. \quad (2)$$

The existence of a nontrivial representation of (2) can then be shown simply by defining the A_i 's, $\langle W |$ and $|V \rangle$ exactly as in ref. [9]. We use here a slightly different notation and define the matrix A_{τ} ($\tau = 1, 2, \dots, m$) by

$$A_{\tau} = |\tau\rangle\langle V| + \sum_{\{\tau_2\}} |\tau\tau_2\rangle\langle\tau_2| + \sum_{\{\tau_2\tau_3\}} |\tau\tau_2\tau_3\rangle\langle\tau_2\tau_3| + \dots \quad (3)$$

The vacuum vector is denoted by $|V\rangle$. This definition implies

$$A_{\tau_1} A_{\tau_2} \dots A_{\tau_L} |V\rangle = |\tau_1 \tau_2 \dots \tau_L\rangle. \quad (4)$$

One can then choose $\langle W |$ such that (2) holds by definition.

Since $|p_0\rangle_{\leftarrow}$ is defined as the stationary state one has $T_{\leftarrow}|p_0\rangle_{\leftarrow} = |p_0\rangle_{\leftarrow}$. The simplest mechanism to achieve this is to imagine that $\mathcal{R}$ produces a 'defect' $\bar{A}$ which is transported through the sequential action of $\mathcal{T}$ to the left end to the chain where it disappears by applying $\mathcal{L}$. This mechanism reads then:

$$\mathcal{T}[A \otimes \bar{A}] = \bar{A} \otimes A, \quad (5)$$

$$\mathcal{R}[A]|V\rangle = \bar{A}|V\rangle, \quad (6)$$

$$\langle W|\mathcal{L}[\bar{A}] = \langle W|A. \quad (7)$$

By defining

$$\bar{A} \otimes A^{\otimes L-1} |V\rangle = \prod_{i=1}^{L-1} T_i R A^{\otimes L} |V\rangle \quad (8)$$

with

$$R = \mathbf{1} \otimes \dots \otimes \mathbf{1} \otimes \mathcal{R}, \quad (9)$$

$$T_i = \mathbf{1} \otimes \mathbf{1} \dots \otimes \mathcal{T}_i \otimes \mathbf{1} \dots \otimes \mathbf{1}, \quad (10)$$

it can be shown analogously to [9] that in fact the defined $\bar{A}$ fulfills the algebra (5)-(7).

This means that in order to solve such a model exactly, the problem of solving the master equation has been reformulated in terms of finding representations of the algebra. Up to now, explicit representations of this algebra are only known for the asymmetric exclusion model ([5],[6],[7]).

We are now able to write down the stationary states for the updates $T_{\rightarrow}$ and $T_{\parallel}$:

$$|P_0\rangle_{\rightarrow} = \langle W | \bar{A}^{\otimes L} |V\rangle, \quad (11)$$

and

$$|P_0\rangle_{\parallel} = \langle W | \bar{A} \otimes A \otimes \bar{A} \otimes A \otimes \dots \otimes \bar{A} \otimes A |V\rangle. \quad (12)$$

By applying $T_{\rightarrow}$ on (11), $T_{\parallel}$ on (12) and using the algebra (5)-(7) one can confirm that these states are in fact the desired stationary states. One still has to confirm that these expressions are not equal to zero. To see this we define

$$C = \sum_{i=1}^m A_i \quad ; \quad \bar{C} = \sum_{i=1}^m \bar{A}_i. \quad (13)$$

The matrices $\mathcal{T}, \mathcal{R}, \mathcal{L}$ have the property that their columns add up to one since the update of a local state has to result in some other local state. Adding up each equation (5)-(7) one obtains

$$[C, \bar{C}] = 0, \quad (14)$$

$$\bar{C}|V\rangle = C|V\rangle, \quad (15)$$

$$\langle W | \bar{C} = \langle W | C. \quad (16)$$

This implies immediately that the norm $\langle W | C^L |V\rangle$ of $|p_0\rangle_{\leftarrow}$ is equal to the norms of $|p_0\rangle_{\rightarrow}$ and $|p_0\rangle_{\parallel}$ which shows that the latter states are nontrivial.

One can now compare the densities of the stationary states produced by the different types of update. Using the formulas (2), (11), (12) and (14)-(16) one gets for the densities $\rho_i(x)$, which are defined as the probability to find a particle of type $\tau = 1, \dots, m$ at location $i = 1, \dots, L$ in the stationary state:

$$\rho_{\tau}(i)_{\parallel} = \begin{cases} \rho_{\tau}(i)_{\rightarrow} & : \quad i \text{ odd} \\ \rho_{\tau}(i)_{\leftarrow} & : \quad i \text{ even} \end{cases}. \quad (17)$$

Note that reversing the order of the the two time steps in the sublattice parallel update would result in reverse arrows in (17).

In order to get similar expressions for the correlation functions we now make use of the identity

$$|V\rangle\langle V| + \sum_{l=1,2,\dots} \sum_{\{\tau_1, \tau_2, \dots, \tau_l\}} |\tau_1, \tau_2, \dots, \tau_l\rangle\langle \tau_1, \tau_2, \dots, \tau_l| = \mathbf{1}, \quad (18)$$

which expresses the completeness of our Fock space.

This can be rewritten as

$$\sum_{l=0,1,2,\dots} C^l |V\rangle\langle V| (C^l)^\dagger = \mathbf{1} \quad . \quad (19)$$

One can now multiply both sides of (19) with $\bar{C}$ and make use of (14) and (15) which leads to

$$C = \bar{C}. \quad (20)$$

This result can actually help to construct explicit representations of the algebra since it reduces the number of matrices $2m$ which have to be represented by one. It also reduces the m^2 equations for the bulk algebra by one and the $2m$ equations for the boundary algebra by two, which can be of significant help.

Note that in the case of random sequential update the same considerations would lead to the result $\bar{C} = 0$, which is fulfilled by the matrix product solution of Derrida et al of the asymmetric exclusion model [3].

One further obtains a relationship between the correlation functions in the stationary states which reads for the two-point correlation function (generalizations to n -point correlation functions are obvious):

$$\langle \tau_i \tau_j \rangle_{\parallel} = \begin{cases} \langle \tau_i \tau_j \rangle_{\rightarrow} & : i, j \text{ odd} \\ \langle \tau_i \tau_j \rangle_{\leftarrow} & : i, j \text{ even} \end{cases} \quad (21)$$

The correlations between odd-even and even-odd sites cannot be expressed so easily. Since the matrix C acts like a transfer matrix between the points of the correlation functions, the relevant length scales are determined by C [5] and will therefore be equal for all three updates.

Since the time evolution of a system can be written as $|p_t\rangle = T^t |p_{t=0}\rangle$ it is useful to ask for the general eigenvectors $|p^E\rangle$ with eigenvalue E of T . It turns out that these can also be constructed ([10],[8]) by writing $E = \epsilon^{L-1} \epsilon_l \epsilon_r$ and multiplying the r.h.s of (8) with $\frac{1}{\epsilon^{L-1} \epsilon_r}$. Details will be given in [8]. It turns also out that by setting $\epsilon = 1$ and $\epsilon_l = \frac{1}{\epsilon_r}$ one has the freedom to substitute (20) with $\bar{C} = \frac{1}{\epsilon_r} C$.

In this letter we have shown that the matrix product formalism can be used to prove the physical equivalence of the sequential and the sublattice parallel update. One gets explicit nontrivial connections between densities and correlation functions in the stationary state.

One can make similar considerations concerning the relation between the stationary states of the random sequential update and the updates discussed in the present work. This work is in progress [8].

Acknowledgments

This work has been performed within the research program of the Sonderforschungsbereich 341 (Köln-Aachen-Jülich). We would like to thank A. Schadschneider and H. Niggemann for fruitful discussions.

References

- [1] Schmittmann B and Zia R K P, *Statistical mechanics of driven diffusive System*, Academic Press, London 1995
- [2] Ligget T, *Interacting particle systems*, Springer, New York 1985
- [3] Derrida B, Evans M R, Hakim V and Pasquier V 1993 *J. Phys. A: Math Gen.* **26** 1493
- [4] Schreckenberg M, Schadschneider A, Nagel K and Ito M 1995 *Phys. Rev. E* **51** 2339
- [5] Hinrichsen H 1996 *J. Phys. A: Math Gen.* **29** 3659
- [6] Rajewsky N, Schadschneider A and Schreckenberg M 1996 *J. Phys. A: Math Gen.* **29** L305
- [7] Honecker A and Peschel I 1996 cond-mat/9606053
- [8] Rajewsky N, Santen L, Schadschneider A and Schreckenberg M, in preparation
- [9] Sandow S and Krebs K 1996 cond-mat/9610029
- [10] Sandow S and Krebs K, private communication